

Submitted to the Astrophysical Journal Letters

X-ray Diagnostics of Grain Depletion in Matter Accreting onto T Tauri Stars

Jeremy J. Drake¹, Paola Testa², Lee Hartmann¹

¹*Smithsonian Astrophysical Observatory, MS-3,
60 Garden Street,
Cambridge, MA 02138*

jdrake@cfa.harvard.edu

²*Kavli Institute for Astrophysics and Space Research, Massachusetts Institute for
Technology,
70 Vassar Street, Cambridge, MA 02139*

ABSTRACT

Recent analysis of high resolution *Chandra* X-ray spectra has shown that the Ne/O abundance ratio is remarkably constant in stellar coronae. Based on this result, we point out the utility of the Ne/O ratio as a discriminant for accretion-related X-rays from T Tauri stars, and for probing the measure of grain-depletion of the accreting material in the inner disk. We apply the Ne/O diagnostic to the classical T Tauri stars BP Tau and TW Hya—the two stars found to date whose X-ray emission appears to originate, at least in part, from accretion activity. We show that TW Hya appears to be accreting material which is significantly depleted in O relative to Ne. In contrast, BP Tau has an Ne/O abundance ratio consistent with that observed for post-T Tauri stars. We interpret this result in terms of the different ages and evolutionary states of the circumstellar disks of these stars. In the young BP Tau disk (age ~ 0.6 Myr) dust is still present near the disk corotation radius and can be ionized and accreted, re-releasing elements depleted onto grains. In the more evolved TW Hya disk (age ~ 10 Myr), evidence points to ongoing coagulation of grains into much larger bodies, and possibly planets, that can resist the drag of inward-migrating gas, and accreting gas is consequently depleted of grain-forming elements.

Subject headings: circumstellar matter—stars: formation—stars: individual (TW Hydrae, BP Tauri)—X-rays: stars

1. Introduction

X-rays, at least in principle, present a powerful method of studying directly the accretion processes of star formation. In the case of classical T Tauri stars (CTTS), the magnetospheric accretion paradigm posits that material is funnelled by magnetic fields onto the star from a truncated disk (e.g. Uchida & Shibata 1984; Bertout et al. 1988), rather than through direct surface interaction with the disk itself. Matter impacts the stellar surface at free-fall velocities of up to a few hundred km s^{-1} . The accreting material is then expected to form a shock at the stellar surface. The resulting shocked plasma temperature is of order a few million K, and it will therefore radiate predominantly in X-rays. For the ranges of mass accretion rates inferred for CTTS of 10^{-6} - $10^{-10} M_{\odot} \text{ yr}^{-1}$ (e.g. Hartigan et al. 1995; Gullbring et al. 1998; Johns-Krull et al. 2000), the X-ray luminosities resulting from accretion should be $\gtrsim 10^{31} \text{ erg cm}^{-2} \text{ s}^{-1}$ —easily sufficient to be observed in nearby associations of T Tauri stars and regions of star formation.

The above scenario of copious accretion-driven X-rays from CTTS contrasts sharply with what is observed. While strong X-ray emission appears to be a ubiquitous characteristic of CTTS (e.g. Feigelson & Montmerle 1999), the source plasma temperatures are an order of magnitude higher than can be produced in accretion shocks and heating can instead be attributed to magnetic processes analogous to coronal activity in late-type main sequence and more evolved stars. Accretion activity is largely revealed by strong UV-optical continuum emission, suggesting that accretion shocks are either formed too deep in the stellar atmosphere to be observed (Drake 2005), or that infall velocities are insufficient to attain X-ray temperatures in the shock. Nevertheless, two stars now stand out as probable exceptions and examples of objects whose high resolution X-ray spectra appear to be produced, at least in part, by accretion: TW Hya (Kastner et al. 2002; Stelzer & Schmitt 2004) and BP Tau (Schmitt et al. 2005).

TW Hya has a dominant plasma temperature of $3 \times 10^6 \text{ K}$, as expected from an accretion shock resulting from free-fall of gas from a truncated disk. Based on the density-sensitive He-like Ne and O lines, both TW Hya and BP Tau seem to be characterised by plasma with high electron densities $n_e \sim 10^{11}$ - 10^{13} cm^{-3} at temperatures of $\sim 3 \times 10^6 \text{ K}$ (Kastner et al. 2002; Stelzer & Schmitt 2004), in contrast to all other single and active binary stars studied in the surveys of Testa et al. (2004) and Ness et al. (2004), where He-like O lines indicate $n_e \sim 10^{10} \text{ cm}^{-3}$. The X-ray spectra of both stars exhibit extremely weak lines of Mg, Si and Fe, and instead are dominated by O and Ne. While this pattern is reminiscent of that seen in very active RS CVn-type binaries (e.g. Huenemoerder et al. 2001), Stelzer & Schmitt (2004) echoed the earlier suggestion of Herczeg et al. (2002) that the metal depletion is instead a signature of the accretion of grain-depleted gas. If the latter is indeed the case, X-rays then

provide a unique means of investigating the matter in the inner accretion disk that is in the process of accreting.

In this *Letter*, we draw on the recent findings of Drake & Testa (2005), based on *Chandra* High Energy Transmission Grating (HETG) spectra of nearby stars, that the Ne/O abundance ratio is remarkably constant in stellar coronae. We use this result to investigate Ne/O in TW Hya and BP Tau, and show that this ratio provides a diagnostic of the depletion of O in grains in the circumstellar disk, and in particular in the very inner disk from which accreting material derives.

2. Observations and Analysis

We use spectral line fluxes for the H-like and He-like resonance lines of Ne and O obtained from *XMM-Newton* observations of TW Hya (acquired 2001 July) and BP Tau (acquired 2004 August) by Stelzer & Schmitt (2004) and Schmitt et al. (2005), respectively; the reader is referred to these works for further details regarding the observations. To provide an additional TW Hya association comparison for TW Hya itself, we also analysed the same lines for the CTTS TWA 5 studied recently by Argiroffi et al. (2005) based on *XMM-Newton* spectra.

TW Hya was observed earlier by the *Chandra* HETG and ACIS-S, on 2000 July 18 at UT10:19 (see Kastner et al. 2002, for further details). Here, we have analysed the products of standard CIAO 3.0 processing, and in particular the Medium Energy Grating spectrum. The measurement of Ne and O line spectral line fluxes and all calculations were performed using the IDL Package for INTeractive Analysis of Line Emission (PINTofALE)¹ following the methods described by Testa et al. (2004). The line fluxes of interest for this study are listed in Table 1.

The conversion of Ne and O fluxes into the Ne/O abundance ratio by number, A_{Ne}/A_O , is described by Drake & Testa (2005). Briefly, an abundance ratio in an optically-thin, collision-dominated plasma can be derived by using lines whose contribution functions $G_{ji}(T)$ —the product of the parent ion population and the line emissivity—have very similar temperature dependence, simply from the ratio of observed line fluxes (corrected for attenuation by interstellar extinction), F_O and F_{Ne} :

$$\frac{A_{Ne}}{A_O} = \left(\frac{G_O}{G_{Ne}} \right) \frac{F_{Ne}}{F_O} \quad (1)$$

¹PINTofALE is freely available from <http://hea-www.harvard.edu/PINTofALE/>

For Ne and O, such a ratio of $G_{ji}(T)$ functions can be constructed from the O VIII, Ne IX and Ne X resonance lines $1s2p\ ^1P_1 \rightarrow 1s^2\ ^1S_0$ and $2p\ ^2P_{3/2,1/2} \rightarrow 1s^2\ ^2S_{1/2}$, combined as follows

$$\overline{\left(\frac{G_O}{G_{Ne}}\right)} = \frac{G_{OVIII}}{G_{NeIX} + 0.15G_{NeX}}. \quad (2)$$

Drake & Testa (2005) found $\overline{G_O/G_{Ne}} = 1.2 \pm 0.1$ for active stars. Ne/O abundance ratios derived for BP Tau, TW Hya and TWA 5 using this method are also listed in Table 1, together with the intervening absorbing columns used to correct the observed line fluxes for attenuation. These Ne/O ratios are compared with those presented by Drake & Testa (2005) for a sample of 21 post-T Tauri stars, including single main-sequence stars, giants, and tidally-interacting binaries in Figure 1.

3. Discussion

Based on the Ne/O ratios found for post-T Tauri stars, Drake & Testa (2005) drew two conclusions: (i) Ne/O is essentially constant in stellar coronae, and in full-disk observations is not susceptible to fractionation effects that often appear to characterise other elements with lower first ionisation potentials (FIPs) (e.g. Drake 2003); (ii) the Ne/O abundance ratio in stars is significantly larger than current assessments of the solar ratio (also illustrated in Figure 1), but is in-line with inference from solar oscillations (Antia & Basu 2005; Bahcall et al. 2005). The constancy of the coronal Ne/O ratio then allows us to use these elements for diagnostics purposes. Firstly, we can determine whether or not the suspected accretion shock has an Ne/O ratio consistent with that of the underlying star, as represented by the constant coronal Ne/O ratio illustrated in Figure 1. A ratio significantly different to the coronal value would provide further evidence that these lines are not formed in a “normal” coronal plasma. Secondly, departures from the coronal Ne/O ratio can then be used to infer compositional peculiarities in the accreting gas.

We conclude from Figure 1 that the Ne/O ratio of $A_{Ne}/A_O \sim 1.0$ we find in TW Hya—in good agreement with the value obtained earlier by Kastner et al. (2002) based on a differential emission measure analysis—is significantly higher than that of more evolved stars (Figure 1). In contrast, Ne/O in BP Tau is perfectly consistent with that found for the rest of the sample of Drake & Testa (2005). This latter result also supports the interpretation of the Ne/O abundance in stellar coronae as representative of the ambient local cosmos: under the assumption that the Ne and O lines in BP Tau are formed from accreting material, as the anomalous O VII line ratios analysed by Schmitt et al. (2005) suggest, the ratio seen in BP Tau indicates that its circumstellar material shares the same Ne/O ratio.

We can confirm that the Ne/O ratio found for TW Hya does not simply reflect an anomalous local environmental composition by comparison with the ratio seen in TWA 5, a CTTS from the same association as TW Hya. The X-ray spectrum of TWA 5 has been analysed recently by Argiroffi et al. (2005): unlike TW Hya and BP Tau, it does not show the high plasma density signature of an accretion shock, and its Ne and O lines can be interpreted as being formed in magnetically-heated coronal plasma. Using the Argiroffi et al. (2005) O and Ne line fluxes (Table 1) we obtain $A_{Ne}/A_O = 0.52 \pm 0.09$ —in good agreement with their result based on a differential emission measure analysis and in excellent agreement with that of the rest of our star sample, except for TW Hya. Similarly, Kastner et al. (2004) find $A_{Ne}/A_O \sim 0.5$ for the TW Hya association multiple, non-accreting, weak-lined T Tauri star system HD 98800 based on *Chandra* HETG spectra.

Herczeg et al. (2002) suggested that the lack of Si in the UV spectrum of the accretion shock of TW Hya was a signature of the accretion of grain-depleted material. Stelzer & Schmitt (2004) invoked their suggestion to explain the metal-poor X-ray spectrum. However, since tidally-interacting binaries have also been shown to exhibit similar metal paucity (see, e.g. reviews by Drake 2003; Audard 2005), it is not immediately clear that the composition of the hot plasma in TW Hya is significantly different. The Ne/O ratio provides this evidence, and supports the conjecture that TW Hya is accreting grain-depleted gas. Unlike comparisons of Ne with metals such as Mg, Si and Fe, the Ne/O diagnostic appears to be robust to the effects of compositional fractionation seen in coronal plasma. Moreover, the Ne and O lines are formed at the temperature of $\sim 3 \times 10^6$ K expected of the accretion shocks on T Tauri stars, whereas dominant lines of Fe XVII and higher charge states, Mg and Si are all formed at higher temperatures.

In the context of shock-produced versus coronal X-rays, it is worth noting that the observed departure from a “normal” chemical composition cannot be mimicked by a corona whose plasma is fed by accretion. The convective turnover time for CTTS is of order a few hundred days (Gilliland 1986; Kim & Demarque 1996) and so accreting plasma, and any abundance peculiarity in these last accumulating fractions, is rapidly subsumed by the underlying star that would feed the corona.

We interpret the different Ne/O ratios in BP Tau and TW Hya as arising from different depletions of O in the accreting gas. The comparison of BP Tau with post-T Tauri stars indicates that it is accreting material which is essentially undepleted. In the case of TW Hya, the O depletion amounts to a factor of ~ 2 or more. Why should BP Tau and TW Hya exhibit distinctly different O depletion in their accreting plasma? We propose that this reflects the different ages and evolutionary states of these stars.

There is some controversy concerning the age of BP Tau, though this is entirely the

result of its uncertain Hipparcos parallax. The Hipparcos data indicate a distance of only 42-70 pc ($\pm 1\sigma$ range)—well in front of the Taurus cloud at 140 pc in which it is generally thought to reside (Wichmann et al. 1998; Favata et al. 1998; Bertout et al. 1999). Based on this distance, Favata et al. (1998) estimated BP Tau to be as old as 35 Myr through comparison with evolutionary tracks. However, both Wichmann et al. (1998) and Bertout et al. (1999) refute the Hipparcos distance based on its low statistical significance. Bertout et al. (1999) find that the astrometric fit to the Hipparcos data for the distance of the Taurus cloud of 140 pc, instead of 42-70 pc, represents only a 2σ deviation, and point out that BP Tau is rather faint for the nominal distance to be reliable. They also show the Hipparcos distance of 32-52 pc for DF Tau—the other Taurus star found by Favata et al. (1998) to be of anomalous proximity—to be highly uncertain owing to binarity and photometric variability. Allowing for a somewhat smaller distance than 140 pc, but not as extreme as the Hipparcos value, Simon et al. (2000) estimate a range of 2-10 Myr. However, all other observational evidence points to BP Tau lying in the Taurus cloud, including radial velocity and proper motion (Hartmann & Stauffer 1989; Bertout et al. 1999), extinction of $A_V \sim 0.5$ (e.g. Gullbring et al. 1996a; Muzerolle et al. 2003, and references therein) and strong interstellar Na I absorption (Gullbring et al. 1996b). Gullbring et al. (1998) estimate the age to be 0.6 Myr.

In contrast, the distance and age of TW Hya are much more secure: with an age of ~ 10 Myr (e.g. Zuckerman & Song 2004), TW Hya appears significantly older than BP Tau. TW Hya is in fact one of the oldest CTTS known to be still accreting. More importantly, near infra-red measurements also show the inner disks of the two stars to be quite different. Veiling measurements for BP Tau show a very large excess of 0.6 at $2.2\mu\text{m}$ (Muzerolle et al. 2003), and $K - L = 0.57 \pm 0.23$ (Kenyon & Hartmann 1995)—typical of the accreting stars in Taurus, 95% of which have $K - L$ colours in the range 0.35-1.2 (Meyer et al. 1997). Muzerolle et al. (2003) find that BP Tau has an optically thick disk whose dust truncation radius is essentially at the corotation radius from which magnetospheric accretion is thought to take place—material from both gas and dust will therefore accrete and this material will not be depleted.

Instead, TW Hya has very little near-infrared excess shortward of $10\mu\text{m}$, with $K - L = 0.25$ (Sitko et al. 2000; Calvet et al. 2002; Uchida et al. 2004). This is more typical of $K - L$ for the *nonaccreting* stars in Taurus that are in the range -0.05 - 0.25 for similar spectral types (K7-M2) (Meyer et al. 1997). Indeed, the inner disk of TW Hya appears to be almost completely cleared (e.g. Calvet et al. 2002; Rettig et al. 2004). Consequently there is very little material in the form of small dust particles available to accrete and replenish the gain-depleted gas. Depletion is also in evidence in the unexpectedly weak or absent Si lines in UV spectra (Valenti et al. 2000; Herczeg et al. 2002), and in low Al and Si abundances in the

gas jet of TW Hya inferred by Lamzin et al. (2004). This depletion is in qualitative accord with the infrared detection of excess emission at $8\text{--}13\mu\text{m}$ attributable to silicates, and the spectral energy distribution that indicates that grain formation and coagulation into larger particles is well advanced (Weinberger et al. 2002; Sitko et al. 2000; Calvet et al. 2002). Larger grains would be expected to settle in the midplane of the circumstellar disk (Chiang et al. 2001; D’Alessio et al. 1999). Calvet et al. (2002) speculate that the truncation of the outer disk at ~ 4 AU inferred from the negligible near-infrared excess could be a developing gap caused by a growing protoplanet. In this scenario, the depletion of metals and oxygen in the X-ray spectrum of the shocked accreting gas would be the result of retention of these elements in bodies of sufficient size that they are not significantly affected by the drag of inwardly migrating gas that feeds the accretion.

4. Conclusions

We have investigated the Ne/O abundance ratios of the accreting gas of the T Tauri stars TW Hya and BP Tau using high resolution X-ray spectra of H-like and He-like resonance lines, and have compared these to the same ratio for the coronal plasma of sample of magnetically-active main-sequence and evolved field stars. In the case of BP Tau, we find an Ne/O abundance ratio in good agreement with that of the coronal plasma of field stars. In contrast, the Ne/O ratio for TW Hya is higher than for the rest of the sample by a factor of two.

We interpret these results in terms of a relative depletion of O in the material accreting from the inner disk of TW Hya as compared with BP Tau. This can be attributed to the different, partially age-related, evolutionary states of the disks of these stars. In the BP Tau disk, dust is still present near the disk corotation radius and can be ionized and accreted, re-releasing elements depleted onto grains. In the more evolved TW Hya disk, evidence points to ongoing coagulation of grains into much larger bodies, and possibly planets, that can resist the drag of inward-migrating gas, and accreting gas is consequently depleted of grain-forming elements.

These results demonstrate the utility of high resolution X-ray spectra of CTTS for probing directly the chemical composition of the accreting gas and the state of evolution of the very inner protoplanetary disk.

We thank the NASA AISRP for providing financial assistance for the development of the PINTofALE package. JJD was supported by NASA contract NAS8-39073 to the *Chandra X-ray Center* during the course of this research. PT was supported by Chandra award

number G03-4005A issued by CXC, and SAO contract SV3-73016 to MIT for support of CXC, which is operated by SAO for and on behalf of NASA under contracts NAS8-39073 and NAS8-03060.

REFERENCES

- Anders, E. & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
- Antia, H. M. & Basu, S. 2005, *ApJ*, 620, L129
- Argiroffi, C., Maggio, A., Peres, G., Stelzer, B., & Neuhäuser, R. 2005, *ArXiv Astrophysics e-prints*
- Asplund, M., Grevesse, N., & Sauval, J. 2004, *ArXiv Astrophysics e-prints*, astro-ph/0410214
- Audard, M. 2005, *Cool Stars, Stellar Systems & The Sun*, 13th Cambridge Workshop, Eds. F. Favata & G. Hussain
- Bahcall, J. N., Basu, S., Pinsonneault, M., & Serenelli, A. M. 2005, *ApJ*, 618, 1049
- Bertout, C., Basri, G., & Bouvier, J. 1988, *ApJ*, 330, 350
- Bertout, C., Robichon, N., & Arenou, F. 1999, *A&A*, 352, 574
- Calvet, N., D’Alessio, P., Hartmann, L., Wilner, D., Walsh, A., & Sitko, M. 2002, *ApJ*, 568, 1008
- Chiang, E. I., Joungh, M. K., Creech-Eakman, M. J., Qi, C., Kessler, J. E., Blake, G. A., & van Dishoeck, E. F. 2001, *ApJ*, 547, 1077
- D’Alessio, P., Calvet, N., Hartmann, L., Lizano, S., & Cantó, J. 1999, *ApJ*, 527, 893
- Drake, J. J. 2003, *Advances in Space Research*, 32, 945
- . 2005, *Cool Stars, Stellar Systems & The Sun*, 13th Cambridge Workshop, Eds. F. Favata & G. Hussain
- Drake, J. J. & Testa, P. 2005, *Nature*, Feb.
- Favata, F., Micela, G., Sciortino, S., & D’Antona, F. 1998, *A&A*, 335, 218
- Feigelson, E. D. & Montmerle, T. 1999, *ARA&A*, 37, 363
- Gilliland, R. L. 1986, *ApJ*, 300, 339

- Grevesse, N. & Sauval, A. J. 1998, *Space Science Reviews*, 85, 161
- Gullbring, E., Barwig, H., Chen, P. S., Gahm, G. F., & Bao, M. X. 1996a, *A&A*, 307, 791
- Gullbring, E., Hartmann, L., Briceno, C., & Calvet, N. 1998, *ApJ*, 492, 323
- Gullbring, E., Petrov, P. P., Ilyin, I., Tuominen, I., Gahm, G. F., & Loden, K. 1996b, *A&A*, 314, 835
- Hartigan, P., Edwards, S., & Ghandour, L. 1995, *ApJ*, 452, 736
- Hartmann, L. & Stauffer, J. R. 1989, *AJ*, 97, 873
- Herczeg, G. J., Linsky, J. L., Valenti, J. A., Johns-Krull, C. M., & Wood, B. E. 2002, *ApJ*, 572, 310
- Huenemoerder, D. P., Canizares, C. R., & Schulz, N. S. 2001, *ApJ*, 559, 1135
- Johns-Krull, C. M., Valenti, J. A., & Linsky, J. L. 2000, *ApJ*, 539, 815
- Kastner, J. H., Huenemoerder, D. P., Schulz, N. S., Canizares, C. R., Li, J., & Weintraub, D. A. 2004, *ApJ*, 605, L49
- Kastner, J. H., Huenemoerder, D. P., Schulz, N. S., Canizares, C. R., & Weintraub, D. A. 2002, *ApJ*, 567, 434
- Kenyon, S. J. & Hartmann, L. 1995, *ApJS*, 101, 117
- Kim, Y. & Demarque, P. 1996, *ApJ*, 457, 340
- Lamzin, S. A., Kravtsova, A. S., Romanova, M. M., & Batalha, C. 2004, *Astronomy Letters*, 30, 413
- Meyer, M. R., Calvet, N., & Hillenbrand, L. A. 1997, *AJ*, 114, 288
- Muzerolle, J., Calvet, N., Hartmann, L., & D’Alessio, P. 2003, *ApJ*, 597, L149
- Ness, J.-U., Güdel, M., Schmitt, J. H. M. M., Audard, M., & Telleschi, A. 2004, *A&A*, 427, 667
- Rettig, T. W., Haywood, J., Simon, T., Brittain, S. D., & Gibb, E. 2004, *ApJ*, 616, L163
- Sanz-Forcada, J., Favata, F., & Micela, G. 2004, *A&A*, 416, 281
- Schmitt, J. H. M. M., Robrade, J., Ness, J.-U., Favata, F., & Stelzer, B. 2005, *A&A*, 432, L35

- Simon, M., Dutrey, A., & Guilloteau, S. 2000, *ApJ*, 545, 1034
- Sitko, M. L., Lynch, D. K., & Russell, R. W. 2000, *AJ*, 120, 2609
- Stelzer, B. & Schmitt, J. H. M. M. 2004, *A&A*, 418, 687
- Testa, P., Drake, J. J., & Peres, G. 2004, *ApJ*, 617, 508
- Uchida, K. I., Calvet, N., Hartmann, L., Kemper, F., Forrest, W. J., Watson, D. M., D’Alessio, P., Chen, C. H., Furlan, E., Sargent, B., Brandl, B. R., Herter, T. L., Morris, P., Myers, P. C., Najita, J., Sloan, G. C., Barry, D. J., Green, J., Keller, L. D., & Hall, P. 2004, *ApJS*, 154, 439
- Uchida, Y. & Shibata, K. 1984, *PASJ*, 36, 105
- Valenti, J. A., Johns-Krull, C. M., & Linsky, J. L. 2000, *ApJS*, 129, 399
- Weinberger, A. J., Becklin, E. E., Schneider, G., Chiang, E. I., Lowrance, P. J., Silverstone, M., Zuckerman, B., Hines, D. C., & Smith, B. A. 2002, *ApJ*, 566, 409
- Wichmann, R., Bastian, U., Krautter, J., Jankovics, I., & Rucinski, S. M. 1998, *MNRAS*, 301, L39+
- Zuckerman, B. & Song, I. 2004, *ARA&A*, 42, 685

d




Fig. 1.— Derived Ne/O abundance ratios by number as a function of the coronal activity index L_X/L_{bol} . Data points represent the A_{Ne}/A_O for the objects in the Drake & Testa (2005) sample, together with TW Hya, BP Tau and TWA 5 analysed here. Error bars are based on the quadrature addition of the uncertainties of line flux measurement. The other hollow symbols correspond to the stars Procyon (F5 IV) and ϵ Eri (K2 V) for which the abundances were taken from the literature (Sanz-Forcada et al. 2004) to better represent the lower ranges of coronal activity. The error-weighted mean Ne/O abundance ratio obtained is $A_{Ne}/A_O = 0.41$, or 2.7 times the currently assessed value (Asplund et al. 2004) which is illustrated by the dotted horizontal line. The recommended values from earlier studies in common usage (Grevesse & Sauval 1998; Anders & Grevesse 1989) are also illustrated.

Table 1. Spectral line fluxes and derived Ne/O abundance ratios (by number).

Source	Spec. Type	d [pc]	L_X/L_{bol}	N_H [10^{20} cm^{-2}]	Fluxes [$10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$] ^e			A_{Ne}/A_O
					Ne IX	Ne X	O VIII	
TW Hya	K7	56	−3.05	2.0 ^b	2.29 ± 0.19^a	1.52 ± 0.14^a	2.9 ± 0.4^a	0.87 ± 0.13
	...	...	...	2.0 ^b	3.64 ± 0.32^b	1.09 ± 0.25^b	3.6 ± 0.3^b	1.06 ± 0.13
BP Tau	K5	140	−3.59	10-20 ^c	0.108 ± 0.029^c	0.29 ± 0.04^c	0.28 ± 0.03^c	0.54 ± 0.12
TWA 5	M1.5	50	−3.1	3.0 ^d	0.59 ± 0.13^d	1.28 ± 0.15^d	1.51 ± 0.9^d	0.52

^aBased on *Chandra* MEG spectra (this work).

^bFrom Stelzer & Schmitt (2004) based on XMM-Newton RGS spectra.

^cFrom Schmitt et al. (2005) based on XMM-Newton RGS spectra.

^dFrom Argiroffi et al. (2005) based on XMM-Newton RGS spectra.

^eFluxes have been corrected for the listed intervening H column densities from Stelzer & Schmitt (2004) for TW Hya, Schmitt et al. (2005) for BP Tau, and Argiroffi et al. (2005) for TWA 5.